


Zambia

Epidemiological Fact Sheet

on HIV/AIDS
and sexually
transmitted
infections



2000 Update




**World Health
Organization**

2 – Zambia

Country Information

Population pyramid, 1999



UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global Surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, guides respective activities. The primary objective of the working group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the working group collaborates closely with national AIDS programmes and a number of national and international experts and institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional and global levels. The Epidemiological Fact Sheets are one of the products of this close and fruitful collaboration across the globe.

The working group and its partners have established a framework standardizing the collection of data deemed important for a thorough understanding of the current status and trends of the epidemic, as well as patterns of risk and vulnerability in the population. Within this framework, the Fact Sheets collate the most recent country-specific data on HIV/AIDS prevalence and incidence, together with information on behaviours (e.g. casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed-upon indicators was not available for many countries in 1999. However, these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The Fact Sheets may also be instrumental in identifying potential partners when planning and implementing improved surveillance systems.

The fact sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the working group would like to encourage all programme managers as well as national and international experts to communicate additional information to the working group whenever such information becomes available. The working group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

Indicators	Year	Estimate	Source
Total Population (thousands)	1999	8,976	UNPOP
Population Aged 15-49 (thousands)	1999	4,102	UNPOP
Annual Population Growth	1990-1998	2.4	UNPOP
% of Population Urbanized	1998	43	UNPOP
Average Annual Growth Rate of Urban Population	1990-1998	2.7	UNPOP
GNP Per Capita (US\$)	1997	370	World Bank
GNP Per Capita Average Annual Growth Rate	1996-1997	1.8	World Bank
Human Development Index Rank (HDI)	1999	151	UNDP
% Population Economic Active			
Unemployment Rate			
Total Adult Literacy Rate	1995	78	UNESCO
Adult Male Literacy Rate	1995	86	UNESCO
Adult Female Literacy Rate	1995	71	UNESCO
Male Secondary School Enrollment Ratio	1996	35.1	UNESCO
Female Secondary School Enrollment Ratio	1996	22.2	UNESCO
Crude Birth Rate (births per 1,000 pop.)	1999	42	UNPOP
Crude Death Rate (deaths per 1,000 pop.)	1999	20	UNPOP
Maternal Mortality Rate (per 100,000 live births)	1990	940	WHO
Life Expectancy at Birth	1998	40	UNPOP
Total Fertility Rate	1998	5.5	UNPOP
Infant Mortality Rate (per 1,000 live births)	1999	80	UNICEF/UNPOP

Contact address:
 UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance
 20, Avenue Appia
 CH-1211 Geneva 27
 Switzerland
 Fax: +41 22 791 4878
 email: surveillance@UNAIDS.org
<http://www.who.ch/emc/diseases/hiv>
<http://www.unaids.org>

Estimated number of people living with HIV/AIDS

In 1999 and during the first quarter of 2000, UNAIDS and WHO worked closely with national governments and research institutions to recalculate current estimates on people living with HIV/AIDS. These calculations are based on the previously published estimates for 1997 and recent trends in HIV/AIDS surveillance in various populations. A methodology developed in collaboration with an international group of experts was used to calculate the new estimates on prevalence and incidence of HIV and AIDS deaths, as well as the number of children infected through mother-to-child transmission of HIV. Different approaches were used to estimate HIV prevalence in countries with low-level, concentrated or generalized epidemics. The current estimates do not claim to be an exact count of infections. Rather, they use a methodology that has thus far proved accurate in producing estimates that give a good indication of the magnitude of the epidemic in individual countries. However, these estimates are constantly being revised as countries improve their surveillance systems and collect more information.

Adults in this report are defined as women and men aged 15 to 49. This age range covers people in their most sexually active years. While the risk of HIV infection obviously continues beyond the age of 50, the vast majority of those who engage in substantial risk behaviours are likely to be infected by this age. The 15 to 49 age range was used as the denominator in calculating adult HIV prevalence.

□ Estimated number of adults and children living with HIV/AIDS, end of 1999

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS, alive at the end of 1999

Adults and children	870000		
Adults (15-49)	830000	Adult rate (%)	19.95
Women (15-49)	450000		
Children (0-14)	40000		

□ Estimated number of deaths due to AIDS

Estimated number of adults and children who died of AIDS during 1999:

Deaths in 1999	99000
-----------------------	--------------

□ Estimated number of orphans

Estimated number of children who have lost their mother or both parents to AIDS (while they were under the age of 15) since the beginning of the epidemic:

Cumulative orphans	650000
---------------------------	---------------

Estimated number of children who have lost their mother or both parents to AIDS and who were alive and under age 15 at the end of 1999:

Current living orphans	447114
-------------------------------	---------------

Assessment of epidemiological situation – Zambia

HIV seroprevalence information among antenatal clinic attendees is available since the mid-1980s from Zambia. In Zambia, Lusaka and Ndola are the major urban areas. HIV prevalence among antenatal women tested in the major urban areas increased from 5 percent in 1985 to 27 percent in 1992 and has remained stable at that rate through 1998. Although overall HIV prevalence rates have remained the same between 1992 and 1998, HIV prevalence among the youngest age group has declined. In 1993, 27 percent of antenatal clinic women less than 20 years of age tested were HIV positive. By 1998, that rate had declined to 17 percent. In 1994 and 1998, HIV prevalence among antenatal clinic women outside the major urban centers remained stable at 14 percent. HIV prevalence ranged from 5 percent to 31 percent of women tested in 18 sites in 1998. In 1994, 14 percent of antenatal clinic women less than 20 years of age who were tested outside of the major urban areas were HIV positive. This rate declined to 6 percent in 1998.

In 1991, 60 percent of male STD patients and 69 percent of female STD clinic patients tested in Lusaka were HIV positive. Outside of Lusaka, 41 percent of female STD patients tested were HIV positive.

4 - Zambia

HIV sentinel surveillance

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV data base maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences is compiled. To provide for a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study on which the medians were calculated are printed at the end of this fact sheet.



The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and – where applicable – other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.

□ HIV prevalence in selected populations in percent (for blood donors: 1/100 000)

Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Pregnant women	Major Urban Areas	N-sites		2		1			1		3	4	4		1		4	
		Minimum		2		11.6			24.5		22.6	22	21.7		26.1		25.9	
		Median		5		11.6			24.5		27	25.15	26.6		26.1		27	
		Maximum		8		11.6			24.5		29.7	27.1	35.3		26.1		29.1	
Pregnant women	Outside Major Urban Areas	N-sites							4	2	6	8	22				18	
		Minimum							9	7.5	5.9	4.9	5				5.2	
		Median							16.5	18	9.95	12.5	13.8				13.9	
		Maximum							30	28.5	23.3	19.2	31.9				31	
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Sex workers	Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																
Sex workers	Outside Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Injecting drug users	Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																
Injecting drug users	Outside Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
STI patients	Major Urban Areas	N-sites		2					2	2	1							
		Minimum		17.9					54	59.7	58							
		Median		22.4					56.5	64.2	58							
		Maximum		26.9					59	68.7	58							
STI patients	Outside Major Urban Areas	N-sites								12								
		Minimum								33.3								
		Median								40.95								
		Maximum								71.4								
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Blood Donors	National	N-sites																
		Minimum																
		Median																
		Maximum																
Blood Donors	Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																
Group	Area		1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Men having sex with men	Major Urban Areas	N-sites																
		Minimum																
		Median																
		Maximum																

Maps of HIV sentinel sites

Mapping the geographical distribution of HIV sentinel sites for different population groups may assist interpreting both the national coverage of the HIV surveillance system and explaining differences in levels and trends of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the UNICEF/WHO HealthMap Programme, has produced maps showing the location and HIV prevalence of HIV sentinel sites in relation to population density, major urban areas and communication routes. Maps illustrate separately the most recent results from HIV sentinel surveillance in pregnant women and in sub-populations at higher risk of HIV infection.



The boundaries and names shown and the designations used on these maps do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement. WHO 2000, all rights reserved.

6 - Zambia

Reported AIDS cases

AIDS cases by year of reporting

1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	Total	Unkn
0	0	0	0	0	1	3	1584	3862	4477	4638	4702	5264	3376	2894	1963	5950	4552	1676			44942	

Date of last report: 31/Jul/97

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases is aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections.

Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of HAART (Highly Active Anti-Retroviral Therapy).

Aids cases by age and sex

Sex	Age	<96	1996	1997	1998	1999	Unkn.	Total	%
All	All								
	0-4								
	5-9								
	10-14								
	15-19								
	20-24								
	25-29								
	30-34								
	35-39								
	40-44								
	45-49								
	50-54								
	55-59								
	60+								
	NS								
Male	All								
	0-4								
	5-9								
	10-14								
	15-19								
	20-24								
	25-29								
	30-34								
	35-39								
	40-44								
	45-49								
	50-54								
	55-59								
	60+								
	NS								
Female	All								
	0-4								
	5-9								
	10-14								
	15-19								
	20-24								
	25-29								
	30-34								
	35-39								
	40-44								
	45-49								
	50-54								
	55-59								
	60+								
	NS								

AIDS cases by mode of transmission

Hetero: Heterosexual contacts.

Homo/Bi: Homosexual contacts between men.

IDU: Injecting drug use. This transmission category also includes cases in which other high-risk behaviours were reported, in addition to injection of drugs.

Blood: Blood and blood products.

Perinatal: Vertical transmission during pregnancy, birth or breastfeeding.

NS: Not specified/unknown.

Sex	Trans. Group	<96	1996	1997	1998	1999	Unkn	Total	%
All	Total	3019							
	Hetero	2794							
	Homo/Bi	19							
	IDU	19							
	Blood	187							
	Perinatal	0							
	Other Known	0							
	Unknown	0							
Male	Total	0							
	Hetero	0							
	Homo/Bi	0							
	IDU	0							
	Blood	0							
	Perinatal	0							
	Other Known	0							
	Unknown	0							
Female	Total	0							
	Hetero	0							
	IDU	0							
	Blood	0							
	Perinatal	0							
	Other Known	0							
	Unknown	0							
NS	Total	3019							
	Hetero	2794							
	IDU	19							
	Blood	19							
	Perinatal	187							
	Other Known	0							
	Unknown	0							

Curable Sexually Transmitted Infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STI are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STI facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Significant also is the observation of a sharp decline in the concentration of HIV in the genital secretions when the infection is treated. Monitoring trends in STI can provide valuable information on the sexual transmission of HIV as well as the impact of behavioural interventions, such as promotion of condom use.

Clinical services offering STI care are an important access point for people at high risk for both AIDS and STI, not only for diagnosis and treatment but also for information and education. Therefore, control and prevention of STI have been recognized as a major strategy in the prevention of HIV infection and ultimately AIDS. One of the cornerstones of STI control is adequate management of patients with symptomatic STIs. This includes diagnosis, treatment and individual health education and counselling on disease prevention and partner notification. Consequently, monitoring different components of STI control can also provide information on HIV prevention within a country.

☐ Estimated incidence and prevalence of curable STIs

STI's	Year	Incidence			Year	Prevalence		
		Male	Female	All		Male	Female	All
Chlamydia trach.								
Gonorrhoea								
Syphilis								
Trichomonas								

Comments:

Sources:

☐ STI Incidence, men

Prevention Indicator 9: Proportion of men aged 15-49 years who reported episodes of urethritis in the last 12 months.

Year	Area	Age	Rate	N=
1996	All		8.9	1732

Comments:

Sources: ZDHS

☐ STI Prevalence, women

Prevention Indicator 8: Proportion of pregnant women aged 15-24 years attending antenatal clinics whose blood has been screened with positive serology for syphilis.

Year	Area	Age	Rate	N=
1997	All		12.0	66000

Comments:

Sources: NASTLP

☐ STI Case management (counselled)

Prevention Indicator 7: Proportion of people presenting with STD or for STD care in health facilities who received basic advice on condoms and on partner notification.

Year	Area	Age	Rate	N=
1996	All	(Male)	66.0	
1996	All	(Female)	91.0	

Comments:

Sources: ZDHS

☐ STI Case management (treatments)

Prevention Indicator 6: Proportion of people presenting with STD in health facilities assessed and treated in an appropriate way (according to national standards).

Year	Area	Age	Rate	N=
1996	All	(Male)	90.0	
1996	All	(Female)	98.0	

Comments:

Sources: ZDHS

8 – Zambia

Health service indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. These efforts may range from reaching out to vulnerable communities through large-scale educational campaigns or interpersonal communication; provision of treatment for STIs; distribution of condoms and needles; creating and enabling environment to reduce risky behaviour; providing access to voluntary testing and counselling; home or institutional care for persons with symptomatic HIV infection; and preventing perinatal transmission and transmission through infected needles or blood in health care settings. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators – such as the percentage of a population with access to health care services; the percentage of women covered by antenatal care; or the percentage of immunized children – may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS – related issues.

☐ Access to health care

Indicators	Year	Estimate	Source
% of population with access to health services – total:	1996	52	ZDHS
% of population with access to health services – urban:	1996	34	ZDHS
% of population with access to health services – rural:	1996	14	ZDHS
Contraceptive prevalence rate (%):	1990-1999	26	UNICEF/UNPOP
% of births attended by trained health personnel:	1990-1999	47	UNICEF
% of 1-yr-old children fully immunized – DPT:	1995-1998	70	UNICEF
% of 1-yr-old children fully immunized – Polio:	1995-1998	70	UNICEF
% of 1-yr-old children fully immunized – Measles:	1995-1998	69	UNICEF
Proportion of blood donations tested:			
% of ANC clinics where HIV testing is available:			
HIV/AIDS Hospital Occupancy Rate (Days):			

Male and female condoms are the only technology available that can prevent sexual transmission of HIV and other STIs. Persons exposing themselves to the risk of sexual transmission of HIV should have consistent access to high quality condoms. AIDS Programmes implement activities to increase both availability of and access to condoms. The two condom availability indicators below are intended to highlight areas of strength and weakness at the beginning and end of the distribution system so that programmatic resources can be directed appropriately to problem areas.

☐ Condom availability (central level)

Prevention Indicator 2: Availability of condoms in the country over the last 12 months (central level).

Year	Area	N	Rate
1997	All	4400	

Comments:

Sources: Medical stores, Care International, PPAZ and Society for Family Health

☐ Condom availability (peripheral level)

Prevention Indicator 3: Proportion of people who can acquire a condom (peripheral level).

Year	Area	N	Rate
------	------	---	------

Comments:

Sources:

Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, intravenous drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of regular behavioural surveys in order to monitor trends in behaviours and target interventions.

☐ **Knowledge of HIV- related preventive practices**

Prevention Indicator 1: Proportion of people citing at least two acceptable ways of protection from HIV infection.

Year	Area	Age Group	Male	Female	All
1996	All	15-19	26.0	25.0	25.5
1996	All	15-49	94.0	100.0	97.0
1996	All	20-24	24.0	23.0	23.5
1996	All	25-49	50.0	52.0	51.0

Comments:

Sources: ZDHS

☐ **Reported non-regular sexual partnerships**

Prevention Indicator 4: Proportion of sexually active people having at least one sex partner other than a regular partner in the last 12 months.

Year	Area	Age Group	Male	Female	All
1990	All	15-19	26.0	13.0	
1995	All	15-19	24.0	8.0	
1990	All	15-49	33.0	8.0	
1995	All	15-49	27.0	5.0	
1990	All	20-24	54.0	11.0	
1995	All	20-24	36.0	7.0	
1990	All	25-39	36.0	6.0	
1995	All	25-39	30.0	3.0	
1990	All	40-49	21.0	4.0	
1995	All	40-49	17.0	2.0	

Comments:

Sources: 1990-Lusak KABP/Behavioural Studies – GPA, 1993. Dynamics and determinants of the HIV epidemic: A review of Zambian epidemiological, demographic and behavioural observations. K, Fylkesnes, Rosemary Mubanga Musonda, Kelvin Kasumba, Zacchaeus Ndhlovu, 1998, 1997
1995-Lusak Dynamics and determinants of the HIV epidemic: A review of Zambian epidemiological, demographic and behavioural observations. K, Fylkesnes, Rosemary Mubanga Musonda, Kelvin Kasumba, Zacchaeus

☐ **Reported condom use in risk sex (gen pop)**

Prevention Indicator 5: Proportion of people reporting the use of a condom during the most recent intercourse of risk.

Year	Area	Age Group	Male	Female	All
1996	All	15-19	68.4	49.7	53.3
1996	All	20-24	35.0	21.8	24.1
1996	All	25-49	10.9	14.6	13.9
1996	All	15-49	26.1	22.0	23.8

Comments:

Sources: ZDHS

10 – Zambia

Knowledge and behaviour

☐ Ever use of condom

Percentage of people who ever used a condom.

Year	Area	Age Group	Male	Female	All
1992	All	15-19		5.6	
1996	All	15-19	23.9	11.7	
1992	All	20-24		15.1	
1996	All	20-24	51.5	24.9	
1992	All	25-29		13.7	
1996	All	25-29	57.3	25.9	
1992	All	30-34		9.8	
1996	All	30-34	45.4	17.2	
1992	All	35-39		6.1	
1996	All	35-39	42.9	12.8	
1992	All	40-44		2.9	
1996	All	40-44	29.5	7.1	
1992	All	45-49		1.6	
1996	All	45-49	18.9	3.7	
1992	All	Total		9.1	
1996	All	Total	38.3	17.0	

Comments:

Sources: Demographic and Health Survey

☐ Median age at first sexual experience

Median age of people at which they first had sexual intercourse.

Year	Area	Age Group	Male	Female	All
1992	All	20-24		16.9	
1992	All	45-49		16.6	
1996	All	20-24		16.0	
1992	All	25-49		16.6	
1996	All	45-49	18.6	16.3	

Comments:

Sources: ZDHS

☐ Adolescent pregnancy

Percentage of teenagers 15-19 who are mothers or pregnant with their first child

Year	Area	Age Group	N	Rate
1992	All	15	384	5.3
1996	All	15	398	4.5
1992	All	16	427	14.7
1996	All	16	419	15.3
1992	All	17	392	29.9
1996	All	17	379	28.3
1992	All	18	380	54.3
1996	All	18	406	46.1
1992	All	19	401	65.6
1996	All	19	401	59.4

Comments:

Sources:

Sources

Data presented in this Epidemiological Fact Sheet come from several different sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations Agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

-
- Duncan, L. E., A. M. Elliott, G. Tembo, et al., 1994, Limitations of the WHO/CDC Clinical Case Definition for AIDS in Africa, *Tropical Doctor*, vol. 24, pp. 11-12.
- Epid. Unit, National AIDS Res. and Control Programme, 1992, Global Programme on AIDS: Sentinel HIV Surveillance on Zambia, World Health Organization, Minister of Health, Zambia, March, unpublished report.
- Fleming, A. F., J. Schneider, I. Wendler, et al., 1987, AIDS in Africa, In: *AIDS The Safety of Blood and Blood Products*, John Wiley and Sons, pp. 241-248.
- Fylkesnes, K., R. Mubanga Musonda, N. P. Luo, et al., 1993, HIV Infection in Pregnant Women in Zambia 1990-93: HIV among Women in Zambia, The Epidemiology and Research Unit, National AIDS/STS/TB and Leprosy Programme, Lusaka, unpublished report, pp. 1-22.
- Fylkesnes, K., R. M. Musonda, K. Kasumba, et al., 1997, The HIV Epidemic in Zambia: Socio-Demographic Prevalence Patterns and Indications of Trends among Childbearing Women, *AIDS*, vol. 11, no. 3, pp. 339-345.
- Fylkesnes, K., R. M. Musonda, K. Kasumba, et al., 1997, Dynamics and Determinants of the HIV Epidemic: A Review of Zambian Epidemiological, Demographic and Behavioural Observations, Paper Presented at the UNAIDS Regional Workshop on "Evidence of Behavioural Change in the Context of HIV Decline in Uganda," 10-13, February, Nairobi, Kenya.
- Fylkesnes, K., Z. Ndhlovu, K. Kasumba, et al., 1998, Studying Dynamics of the HIV Epidemic: Population-Based Data Compared with Sentinel Surveillance in Zambia, *AIDS*, vol. 12, no. 10, pp. 1227-1234.
- Hira, S., G. Bhat, J. Kamanga, et al., 1989, Perinatal Transmission of HIV-1 in Lusaka, Zambia, *British Medical Journal*, vol. 299, no. 6718, pp. 1250-1252.
- Hira, S., J. Kamanga, G. Tembo, et al., 1991, Control Strategies in STD/HIV Clinic in Zambia: a Demonstration Project, VII International Conference on AIDS, Florence, Italy, 6/16-21, Poster W.C.3082.
- Johns Hopkins University, 1986, AIDS -- A Public Health Crisis, Population Information Program, Population Reports, Issues in World Health, July-Aug., Series L, no. 6, pp. 194-228.
- Kanyama, I. D. A., 1991, Sentinel Surveillance of HIV Infection in Northern Zambia, VII International Conference on AIDS, Florence, Italy, 6/16-21, Abstract M.C.3301.
- Melbye, M., E. K. Njelesani, A. Bayley, et al., 1986, Evidence for Heterosexual Transmission and Clinical Manifestations of Human Immunodeficiency Virus Infection and Related. ..., *Lancet*, Nov. 15, vol. 2, no. 8516, pp. 1113-1115.
- Msiska, R., 1992, Overview of the HIV/AIDS Situation in Zambia, Ministry of Health, Zambia, pp. 1-11.
- National HIV Sentinel Surveillance Team, 1999, *Zambian HIV Sentinel Surveillance: Patterns and Trends of the HIV Epidemic*, Ministry of Health, Central Board of Health, June report.
- Sandala, L., P. Lurie, M. R. Sunkutu, et al., 1995, "Dry Sex" and HIV Infection among Women Attending a Sexually Transmitted Diseases Clinic in Lusaka, Zambia, *AIDS*, vol. 9, suppl. 1, S61-S68.
-

Websites: www.aids.africa.com

12 – Zambia

Annex: HIV Surveillance data by site

Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Pregnant women	Major Urban Areas	Lusaka	8		11.6			24.5									
		Lusaka - Chelstone								22.6	26.8	24.7		26.1		25.9	
		Lusaka - Chilenje								27	22	35.3				27.3	
		Lusaka - Kalingalinga									23.5	21.7				26.7	
		Lusaka - Matero								29.7	27.1	28.4				29.1	
		Ndola	2														
Pregnant women	Outside Major Urban Areas	Chikankata								23.3	19.2						
		Chilonga										16.8					
		Chipata										30.3				27.3	
		Chitokoloki										7.1					
		Ibenga										11.4				10.1	
		Isoka										10.6				11.7	
		Kabompo						20				5				9.3	
		Kabwe										29.5				27.2	
		Kalabo								5.9	4.9	10.2				12.8	
		Kamuchanga										23.1					
		Kapiri Mposhi										13				16.5	
		Kasaba										12				5.2	
		Kasama										23.8				14.7	
		Kashikishi										14.6				13	
		Livingstone										31.9				31	
		Macha								7.9	10	9.1				8	
		Mansa										23.6				21.1	
		Minga								12.9	17.7	9.6				10.3	
		Mongu										28.4				27.3	
		Mporo Kosso										12.9					
		Mukinge						13	7.5		9.7	9.5				8.8	
		Mwinilunga						9			9.5						
		Nchelenge								11.4	15.1						
		Ndola														27.5	
		Samfya										20					
		Solwezi						30	28.5	8.5	15	24.2				20.2	
Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Sex workers	Major Urban Areas																
Sex workers	Outside Major Urban Areas																
Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Injecting drug users	Major Urban Areas																
Injecting drug users	Outside Major Urban Areas																
Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
STI patients	Major Urban Areas	Ndola	17.9														
		Lusaka (1)						59									
		Lusaka (2)						54									
		Lusaka (Males)	26.9														
		Lusaka (Females)								58							
		Lusaka Province (Males)							59.7								
		Lusaka Province (Females)							68.7								
STI Patients	Outside Major Urban Areas	Central Province (Females, 1)							33.3								
		Central Province (Females, 2)							53.1								
		Eastern Province (Females, 1)							40								
		Eastern Province (Females, 2)							34.5								

12 (contd) – Zambia

Annex: HIV Surveillance data by site

Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
STI Patients	Outside Major Urban Areas	Luapula Province (Females, 1)							35.3								
		Luapula Province (Females, 2)							41.1								
		Northern Province (Females, 1)							49.3								
		Northern Province (Females, 2)							71.4								
		Southern Province (Females, 1)							40.8								
		Southern Province (Females, 2)							50								
		Western Province (Females, 1)							35.1								
		Western Province (Females, 2)							50								
Group	Area		1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Blood Donors	National																
Blood Donors	Major Urban Areas																
Blood Donors	Outside Major Urban Areas																